

NAME _____
DATE _____
WEIGHT _____ (_____ %)
LENGTH _____ (_____ %)
HEAD _____ (_____ %)



FOR YOUR 2 MONTH OLD

FEEDING

Breastfeed your baby about every 3 hours (8 feedings per day). Feed until your baby seems content. Your baby may “cluster” the feedings in the evening to prepare for longer stretches of nighttime sleep. Continue prenatal vitamins while breastfeeding.

If you breastfeed exclusively, add a vitamin D supplement. 400 IU/day of vitamin D is recommended and is available over the counter at drugstores in many forms (i.e. BabyD drops). Formula fed babies should also be on a vitamin D supplement unless they are drinking 32 ounces a day.

If you are not breastfeeding, feed your baby 4-5 ounces every 3-4 hours for a total of 26-32 ounces per day.

Review the formula preparation instructions carefully. Do not heat formula in the microwave.

Healthy babies do not require extra water.

Do not give honey/Karo syrup in the first year.

Do not offer your baby cereal or other solids until at least 4 months of age. Cereal will not help a baby sleep through the night. Additionally, please do not put cereal in your baby's bottles unless instructed by your healthcare provider for medical purposes.

DEVELOPMENT (2-4 MONTHS)

Raises body when lying on tummy, holds head up, may bear weight on legs.

Smiles, coos, laughs, may blow bubbles, cries change according to needs.

Visually follows any object, responds to sound and voices by quieting or cooing.

Swats, reaches, grabs, may hold a rattle.

SLEEP

Always place your baby on his back to sleep on a firm, flat mattress. Avoid blankets and soft toys in the crib.

The average baby sleeps 15 1/2 hours per day between 1 and 4 months of age. It is normal for your baby to sleep more on some days and less on other days.

By 4 months of age, most babies have longer stretches of sleep at night and 3 daytime naps.

Some parents are concerned about head flattening; tummy time to play will help avoid this.

ELIMINATION

Breastfed babies have yellow stools with a pasty, watery, or curd-like consistency.

Bottle fed babies usually have soft brown, green, or yellow stools.

The frequency varies widely - a few per day to a few per week is normal.

SAFE TOYS

Choose toys that are appropriate for your baby's age.

Check for small or loose parts, like eyes, and ribbons (cut them off). All toy pieces should be no smaller than 1 5/8 inches with no sharp or pointed edges.

Avoid old painted toys that may contain lead.

Do not string any toys across the crib.

FOR YOUR 2 MONTH OLD

SAFETY

Always use a car seat made for an infant and place it rear-facing in the back seat until 2 years of age. Be sure to adjust the shoulder strap as your baby grows.

Always make sure your baby sleeps in a safe place. The crib slats should be less than 2 3/8 inches apart. Bumper pads should not be used.

Take an infant CPR class.

Set your hot water heater to 120° F to prevent burns. Never drink hot liquids while holding your baby.

Never leave your baby unattended on the bed, sofa, changing table, or in the bathtub.

OTHER THINGS TO KEEP IN MIND

Hold, talk, read, and sing to your baby.

Show your baby affection – you cannot spoil a young baby!

Know the signs of illness – temperature **over 100.5** rectally, vomiting (not just spit-up), diarrhea, or failure to eat several feedings in a row. Call our office to see if these symptoms need further evaluation.

Shaking or spanking your baby may cause serious injury or death.

Watch your pets carefully around your baby.

Install and maintain smoke and carbon monoxide detectors.

Do not smoke around your baby, especially inside your home or car. It may lead to respiratory problems and allergies, and can increase the risk for sudden infant death syndrome (SIDS). This is a great time to consider quitting!

Protect from sun exposure with clothing, hats, blankets, or an umbrella. Sunscreen (SPF 15 or higher) may be used when sun exposure is unavoidable, but preferably not on the face or hands.

SUGGESTED READING

“The Nursing Mother Companion”
“What to Expect the First Year”
“Guide to Your Child’s Sleep (AAP)”

WEBSITES

www.carypediatriccenter.com
www.aap.org
www.cdc.gov/vaccines
www.seatcheck.org
www.healthychildren.org

ACETAMINOPHEN DOSAGE (160 mg/5ml) (EVERY 4-6 HOURS)

6-11 lbs = 1.25 ml

12-17 lbs = 2.5ml

No ibuprofen should be given prior to 6 months of age.

TESTS/IMMUNIZATIONS FOR TODAY

Your baby will receive immunizations today. Please review our information sheets or you may visit www.aap.org for vaccine information.

ADDITIONAL INSTRUCTIONS

YOUR NEXT APPOINTMENT IS AT 4 MONTHS

at the four month check-up...

- Your baby will have a physical examination.
 - Your questions and concerns will be answered.
 - Guidelines for feeding solids will be reviewed.
 - Your baby will receive the second set of vaccines.
-